



DARK ENERGY
SURVEY

FNAL-ANL PreCam Reductions

Douglas L. Tucker
(FNAL)

DES Collaboration Meeting
ICG, Portsmouth

PreCam Parallel Session
29 June 2011



DARK ENERGY
SURVEY

Data Processing

- DES-Brazil Effort
 - The official data processing.
 - Uses a PreCam-specific version of the Quick Reduce Pipeline.
 - Quick Reduce in turn uses the DESDM code.
- FNAL/ANL Effort
 - Uses custom scripts in order to understand the data and obtain some quick results.
 - Provides feedback to the official data processing.
 - Most of the *data processing* by Sahar Allam, Douglas Tucker, Kyler Kuehn, and Hope Head, in consultation with Huan Lin, Steve Kuhlmann, Hal Spinka, Tomasz Biesiadzinski, Michael Schubnell, and others.
 - Most of the *data analysis* is being performed at ANL (Kyler, Steve, and Hal), FNAL (Sahar, Huan, Douglas), and UM (Michael). (See Kyler's talk.)



DARK ENERGY
SURVEY

“Golden Nights”

- “Golden Nights”
 - A set of 5 nights with robust FITS headers, no known problems, and target observations in SDSS Stripe 82:

Night	# of Target Fields in Stripe 82				
	g	r	i	z	y
R2010-12-15UT	1	0	40	29	11
R2011-01-07UT	12	0	0	3	0
R2011-01-08UT	0	7	0	10	0
R2011-01-12UT	0	0	10	19	14
R2011-01-17UT	0	0	3	0	0

- Used by both data processing efforts for rapid testing and algorithm development.



DARK ENERGY
SURVEY

FNAL-ANL Processing Methods/Steps (I)

1. A suite of home-grown python scripts are written using (primarily) pyFITS and (occasionally) pyraf.
2. A Master Bias are created by median-combining all good bias frames from entire November – January PreCam observing block.
3. A set of Master Dome Flats are created by median-combining all good flat frames from entire November – January PreCam observing block.
 - Pro: dome flat lamp problems make it difficult to do night-by-night or even week-by-week Master Dome Flats, esp. in late-December and in January.
 - Con: dust specks on the dewar window moved, esp. between PreCam re-mountings.
4. Row-by-row overscan subtraction is performed (takes care of horizontal banding).
5. Horizontal streaking correction is performed on bias-subtracted, flat-fielded science and standard star images. (Important code provided by Tomasz Biesiadzinski and modified by Sahar Allam.)



DARK ENERGY
SURVEY

FNAL-ANL Processing Methods/Steps (II)

6. Illumination/shutter correction maps are created by median-combining processed on-sky images (standard star fields, science targets)
 - One map per filter per exposure time.
 - A night's worth of images? A week's? Kyler Kuehn is investigating this.
7. To simplify analysis, the data for both CCDs are combined into a single FITS image (with a gap in the middle).
8. For later reductions, IRAF fixpix is used to clean bad pixels/columns.
9. Astrometry/WCS keyword values are corrected first by matching against 2MASS (astrometric pre-burner) and then by using IRAF ccmmap routine.
10. Use of SCAMP is being investigated by Michael Shubnell and a summer student.
11. To optimize S/N of fainter stars, PSF photometry (PSFex? DAOPHOT?) will likely need to be used. Hope Head (summer undergrad intern at FNAL) may be investigating this later this summer.




DARK ENERGY
SURVEY

Reduced Data Sets

- FNAL (“v1”)
 - 14 nights processed (superset of Golden Nights)
 - Image de-trending (including horizontal streaking correction), basic astrometric calibration, sextractor catalogs
 - Nearly all analyses to date have been performed on this data set
- FNALv2
 - 49 nights processed (2010-Dec-1 UT → 2011-Jan-18 UT)
 - Just through image processing (no astrometric corrections or sextractor catalogs) so far. Hope Head will be working on astrometry/cataloging.
 - FNAL (“v1”) + IRAF fixpix + horizontal streaking image quality flags in FITS headers
 - Start moving analysis to these reduced data (or to FNALv3?)
- FNALv3
 - Just starting
 - Description: FNALv2 + improved horizontal streaking and image quality flags ⁶




- Experimental Astrophysics Group (EAG) SDSS/DES cluster at Fermilab (e.g., des06.fnal.gov)





DARK ENERGY
SURVEY

End



Credit: S. Allam

A segment of *i*-band PreCam observations in Stripe 82.
FNAL(v1) reductions.
~20 sq deg.




DARK ENERGY
SURVEY

Extra Slides



DARK ENERGY
SURVEY


A Processed *i*-band PreCam Image from Jan 13





DARK ENERGY
SURVEY


Results: Horizontal Banding & Streaking





DARK ENERGY
SURVEY

Results: Horizontal Banding & Streaking






DARK ENERGY
SURVEY


Results: Horizontal Banding & Streaking

A Pretty Bad Case of Banding and Streaking


Original Image



After row-by-row
overscan subtraction



After horizontal
streaking correction







DARK ENERGY
SURVEY

Results: Horizontal Banding & Streaking

Original Image




After horizontal
streaking correction





Dome Flat Lamp Output vs. Time

DARK ENERGY
SURVEY




Credit: Sahar Allam



DARK ENERGY
SURVEY

Results: Horizontal Banding & Streaking

- Horizontal banding & streaking affect $\approx 40\%$ of the raw PreCam standard star field and science target images.
- After correcting, horizontal banding & streaking affect only about 6% of the processed images.




Credit: S. Allam



DARK ENERGY
SURVEY

Results: Initial Photometry for a Single Image



No corrections for:





- overall ZP
- color term
- star flat



Results: Photometry over a Full Night

DARK ENERGY
SURVEY

- Night of 13 Jan 2011 UT.
- All data from that night matching the extended list of USNO $u'g'r'i'z'$ standards.
- Corrections for overall ZPs and for airmass (using site-average first-order extinction coefficients)
- No correction for color terms.
- RMS = 2-4% (mag < 13.0).



Credit: S. Kuhlmann, H. Spinka